

Assessment of Cowpea (*Vigna Unguiculata*(L.) Walp) Germplasm for Agronomic Traits in Seed Production

¹Magashi A. I., ² Gaya A.G., ³ Daraja Y.B.. and ⁴ Isah S.D. ⁵Ado M. ⁶Almu H. ⁷Ahmad D. and ⁸Umar I.

Abstract—A tremendous increase in production of cowpea depends on the introduction of improved varieties and better methods of high quality cowpea seed production, resistance to stress factors, high nutritional value, short vegetation period and high yielding potential. The study was conducted to identify different cowpea varieties which will perform better under specific drought condition with subsequent evaluation of the germplasm for agronomic traits pertinent to seed production. Cowpea germplasm comprising of forty-five (45) genotypes were assessed for agronomic traits in seed production under Gaya conditions, at the KUST experimental field station, during 2013-2015 rainy seasons. The germplasm were obtained from the genes banks and from farmers in Kano and Niger republic respectively. All the forty five germplasm were characterized based on the following parameters: incidence and severity of diseases, agronomic traits as indicators of drought tolerance, grain yield, and earliness and evaluated for days to seedling emergence, germination percent, leaf formation, days to flowering, days to pod formation, number of pods per plant, number of seeds per pod and weight of 100-seeds varied significantly ($P=0.05$) between genotypes. The findings revealed that the less incidence and severity of seed rot, cowpea mosaic, cowpea mottle virus disease, bacterial blight, and anthracnose and nematode infections were recorded in favor of genotypes; KUST-FA-13-35, and KUST-FA-13-41. Genotypes KUST-FA-13-35, KUST-BK-13-4 and KUST-BK-13-5 took between 3-5 days to emerge with 80% percentage emergence and took 14 days from planting to 50% flowering. They have relatively high number of pod per plant number of seeds per pod, highest 100 seed weight with less severity and incidences of diseases. Therefore these varieties are considered to be the best genotypes for seed production under Gaya-Kano State conditions.

Keywords— Cowpea, Germplasm and Agronomic traits, Seed..

¹Magashi A. I., ² Gaya A.G., ³ Daraja Y.B.. and ⁴ Isah S.D. ⁵Ado M. ⁶Almu H. ⁷Ahmad D. and ⁸Umar I. .

^{1,2, 3, 5&8} Department of crop science, Faculty of Agriculture and Agricultural Technology, Kano University of Science and Technology, Wudil, Kano State.

⁴Department of Forestry fisheries and wild life

⁶Department of soil science

⁷Department of Agricultural economics and extension

Corresponding author: Magashi A. I Department of Crop Science, Kano University of Science and Technology, Wudil, Kano State, Nigeria. Email magashi2@yahoo.com. Mobile phone: +2348023627191 and +2348037173580

I. INTRODUCTION

MORE than half of the populations of the people of Kano State are farmers and they practice subsistence farming. The small scale commercial farming is so insignificant that it does not even commensurate with amount of agricultural produce required to the State. There is need to come up with a realistic strategy of boosting sustainable agricultural production, which may not only ensure food security in the State but will enhance the empowerment of common citizen, making them more self-reliant and prosperous. A tremendous increase in food supply at the age of green revolution of mid-20th century resulted in the introduction of better methods of production and improved seed of cultivated crop. [15]

Cowpea (*Vigna unguiculata* L) is the second most important class of food crops grown, providing man with food and feed and is a major source of protein, minerals and vitamins in daily human diets and is equally important as nutritious fodder for livestock among the popular crops grown in Central and West Africa, [1,4 and 17].

Grain yield and its quality are primary breeding objectives of nearly all cowpea breeding programs. The accomplishments of some of these programs have been described by others [11 and 12]. To achieve sustainable crop productivity, maximum yield and quality crop, varieties with potential for high quality cowpea seed need to be identified and consider as an important aspect of cultural practice. Understanding the quantitative characters of varieties for potential seed production will pave way to elucidate the genetic factors responsible for all the identified traits.[3] Consequently, the genes identified can be manipulated to the advantage of farmers.

A lot is known about the extent and structure of genetic variation, and the potential for cowpea crop improvement through domestication, selection and/or breeding of most indigenous cowpea, A number of factors pose as a hindrance to its cultivation. For instance, its relatively low yield is also an impediment to commercial production [2 and 6]

The study was conducted to assess cowpea germplasm for agronomic traits pertinent to seed production a recommend best varieties with resistance to stress factors, high nutritional value, short vegetation period and high yielding potential. Thus enabling cowpea seed producers to receive a boom and solve several problems to come when human population will reach

10 billion in the nearer future

LSD at $P=0.05$.

II. MATERIAL AND METHODS

III. RESULTS AND DISCUSSIONS

A. Genotypes selection

Forty-five (45) genotypes were evaluated for agronomic traits and seed production under KUST Research farm-Gaya conditions in 2014-2015 rainy seasons. The germplasm included 45 genotypes (Table 1), which were obtained from the genes banks and from farmers in Kano and Niger republic (Zinder, Matamaye, Magaria, Mirria, Birnin Konni, Maradi and Gidan Rumji), respectively and a commercial type, was included as a check

B. Experimental Procedure

The genotypes were grown in KUST Farm Plot Gaya, during 2014 and 2015 rainy seasons. Two seeds of each genotype were planted per plot. The varieties were arranged in a completely randomized design (RCBD) with three replicates.

The land was prepared to produce a firm fine seedbed. With emphasis placed on making the bed firm enough to allow for a relatively shallow seed placement. Planted at 2 cm. This was expected to allow moisture to move upward in the soil profile and provides more upward in the soil profile and provides more moisture for the germinating seed. The standard field management practices were strictly observed. Weed spectrum was assessed and an appropriate weed control program was used.

The harvesting begun when two-thirds to three-quarters of the seed pods have turned dark brown or black.

C. Data collection and Analysis

Data on desirable traits was scored based on; incidence and severity of some diseases, and recorded days to seedling emergence, germination percent, leaf accumulation, days to flowering (vegetable production traits), days to pod formation, pods per plant, number of seeds per pod and weight of 100 seeds (seed production traits). Day to flowering was taken when 50% of the plants had flowered while days to pod formation was recorded when 50% of the plants had formed pods. To determine pods per plant, ten plants were randomly selected per genotype in the three replicates and the number of pods in each one of them counted when fully formed and an average number of pods calculated using the data from the ten plants in the three replicates. Similarly for the determination of number of seeds per pod, ten plants were randomly selected in each genotypes for each of the replicates and 5 pods were pick at random from each of the plants and the seeds contained in each pod was counted and recorded and eventually an average per plant calculated. The 100 seed weight was determined by counting 100 seeds of each of the genotypes replicated three times and weighed using a sensitive weighing balance and an average for the 100 seeds calculated. The data collected were subjected to analysis of variance using Genstat statistical package and means were compared using Fisher's protected

A. Incidence and severity of diseases

Incidence and severity of the following disease were observed and recorded see Tables 3, 4, 6, 8, 10, 12 and 19: Seed rot disease, Cowpea mosaic virus, Cowpea mottle carmo-virus, Cowpea aphid-borne mosaic virus, Bacterial blight, anthracnose and Nematode infection. The less incidence and severity of seed rot, cowpea mosaic, cowpea mottle virus disease, bacterial blight, anthracnose and nematode infections were recorded in favor of genotypes; KUST-BK-13-2 and KUST-FA-13-41. (Table 20).

B. Days to seedling emergence

Genotypes differed significantly ($P=0.05$) in the number of days taken from planting to seedling emergence (Table 21). The days to emergence ranged between 3 days 10 days (table 21). Sixteen out of the 45 genotypes evaluated took 3 to 6 days to emerge and only 6 took longer (8-10 days) to emerge compared to others. Temperature and light have been shown to have a significant influence on germination of cowpea, where best conditions for seed germination are 20-30 °C in darkness. Besides, proper processing of seed determines the quality of the seed for example sun-drying of cowpea plant seeds improve the mean germination time, seedling vigour and overall germination percent compared with shade dried seeds. In addition, majority of the genotypes used were obtained from the Gene bank having been stored for a long period and therefore there is need for regeneration of germplasm stored in the gene bank periodically.

Cowpea seeds are negatively photosensitive and the effects of photo inhibition increases at temperatures lower than 20°C and germination is also influenced by physiological maturity of the seeds. The consequence of genotypes that take a long time to germinate may result in a poor final plant stand and hence low yields per unit area. The longer seeds are in the soil before germination or the slower the germination, the greater the chances are for soil disease and insects to attack the seeds. [8 and 10].

C. Germination Percent

There were no significant differences among genotypes for germination percent (Table 21). The germination percent ranged from 10 to 100%. Only two genotypes KUST-BK-13-4 and KUST-BK-13-5 had a germination of 100%, while Genotypes KUST-BK-13-9, KUST-BK-13-10 and KUST-DM-13-11 had germination ranged from 10% - 20%. The number of plants established from a given weight of seed depends on size of seeds and percent of those seeds that are viable and can grow into established plants.

D. Days to 50% Flowering

The number of days to flowering varied significantly ($P=0.05$) among genotypes (Table 21). The shortest duration to flowering

was 11 days for (KUST-BK-13-3, KUST-BK-13-5, KUST-DN-13-12 and KUST-BK-13-2), whereas the longest period to flowering was 18 days. KUST-BK-13-4, KUST-BK-13-6, KUST-DM-13-14 and KUST-MR-13-20.

E. Pods per Plant

The number of pods per plant varied significantly ($P=0.05$) among genotypes (Table 22). The lowest number of pods per plant (2) was recorded in KUST-KD-13-18 and KUST-BK-13-7 and highest number of pods per plant (30) was recorded in KUST-BK-13-10. Majority of genotypes had between 8 and 20 pods per plant. The primary components of seed yield are number of pods per plant and seed weight. Seed yield is a complex character with polygenic inheritance having positive or negative effects on yield component traits. In other crops, seed yield is strongly correlated with number of pods per plant. Taking this into account, genotype with high number of pod per plant would be expected to yield more seed than genotypes that have a lower number of pods per plant. However, this can only be concluded after assessing the seed weight of different genotypes. Salehi et al. reported that the result of stepwise multiple regression analysis based on seed yield as a dependent variable and other traits as independent variables, pods per plant explained 83.2% of the total variation suggesting that the number of pods per plant may be the main factor determining seed yield.

F. Seeds per Pod

There were significant variations of genotypes for the number of seeds per pod (Table 22). The highest number of seeds per pod (11-13 seeds/ pods) was recorded in 17 genotypes, while the lowest number of seeds (7 seeds) per pod was recorded in DAN-YAGAJI. The number of seeds in a pod can vary widely among plant species, individual plants within a species and fruits within a plant. Factors and processes affecting seed production might be achieved with studies that incorporate variation at all these levels.

G. 100-Seed weight

The genotypic effect was significant ($P=0.05$) for 100-seed weight (Table 22). The 100-seed weight is one of the most important criteria in seed quality determination. It determines embryo size and seed storage for germination and emergence. High 100-seed weight increases germination percent and seedling emergence. Thus, seed weight has a large effect on seed germination, seed vigor, seedling establishment and yield production. For commercial purposes, genotypes with a high seed weight will be preferred by farmers as seeds are sold by weight.

G. Grain Yield

Table 23 shows that KUST-FAR-13-42 recorded the highest ($P\leq 0.001$) grain yield among the 45 varieties investigated. However, UST-FA-1329, KUST-FA-13-30, KUST-MG-13-22), KUST-KD-13-188 and KUST-FA-13-40 recorded the

lowest ($P\leq 0.001$) grain yield with no significant difference with some varieties.

IV. CONCLUSION

The less incidence and severity of seed rot, cowpea mosaic, cowpea mottle virus disease, bacterial blight, anthracnose and nematode infections were recorded in favor of genotypes; KUST-BK-13-2, and KUST-FA-13-41. KUST-BK-13-2, KUST-BK-13-4 and KUST-BK-13-5 took between 3 days to emergence, had 80-100 percentage emergence and took 14 days from planting to 50% flowering, with relatively high number of pod per plant number of seeds per pod especially KUST-BK-13-2 (Table) they are considered to be the best genotypes for production under Gaya, Kano, Sudan Savannah. Whereas genotypes; KUST-FA-13-44, KUST-FA-13-41, KUST-FA-13-31, KUST-DM-13-12, KUST-MR-13-19 and KUST-KD-13-17 took more than 24 days from planting to 50% flowering, produced 2-18 pods per plant. Genotypes KUST-BK-13-2 was found to be the best performing under Gaya-Kano State conditions.

REFERENCES

- [1] Alyenemy, M.N (1998). Effect of drought and growth and yield of cowpeas (*Vigna unguiculata* L), Vol. 10 Science (1), 41 – 51
- [2] Ashley, J. (1993). Drought and crop adaptation. Pages 46 – 67 on dryland Farming in Africa, edited by J.R.J Rowland Macmillan Press Ltd, UK.
- [3] Aryeety, A.N and liang, E. (1973). Inheritance of yield components and their Correlation with yield of cowpea (*VignaUnguiculata* (L) Walp) Euphytica 22: 386 – 396
- [4] Barrett, R.P (1990): legumes species as leaf vegetables in: Janick, J, and simon, J.E (Eds). Advance in new crop
- [5] Bates, L.S., R.P. Waldren, and I.O. Teare. 1973. Rapid determination of free proline in water stress studies. Plant and Soil 38: 205.
- [6] Cisse, N, Ndiaye, M. Thiaw, S, Hall, A.E, (1978). Registration of melakli Cowpea crop science 37
- [7] Ceccarelli, S. And S. Grando (1996). Drought as a challenge for the plant Breeder plant growth regulators, 20: 149 – 155
- [8] Dennis R. Decoteau (2005). Principles of plant science Inc. Ch17, PP 295
- [9] Ehlers.J.D, Hall, A.E, (1998). Heat tolerance of contrasting cowpea
- [10] Evans. L.T (1974), the physiological basis of crop yield in evans, L.T. (ed) Crop physiology press 327 – 355
- [11] Hall, A.E, R.G. Mutters, K.T. Hubick, and G.D. Farquhar. 1990. Genotype differences in carbon isotope discrimination by cowpea under wet and dry field conditions. Crop Science 30: 300– 305.
- [12] Hall, A.E., S. Thiaw, A. Ismail, and J.D. Ehlers. 1997. Water-use efficiency and drought adaptation of cowpea. Pages 141–146 in Advances in cowpea research, edited by B.B. Singh, D.R. Mohan
- [13] Krishnamurthy, L.C, C. Johansen, and O. Ito. 1996. Genotypic variation in root system development and its implication for drought resistance in Chickpea. Pages 235–250 in Roots and nitrogen in cropping systems of the semiarid tropics, edited by O. Ito, C. Johansen, J.J. Adu-Gyamfi, K.Katayama, J.V.K. Kumar Rao, and T. J. Rego. JIRCAS and ICRISAT, Hyderabad, India

[14] Morgan, J.M. 1984. Osmoregulation and water stress in higher plants. Annual Review of Plant Physiology 35: 299–319.

[15] Raj, K.E. Dashiell, and L.E.N. Jackai. Copublication of International Institute of Tropical Agriculture (IITA) and Japan International Centre for Agricultural Sciences (JIRCAS). IITA, Ibadan, Nigeria.

[16] Sadras, V.O, and S.P. Milroy (1996). Soil – Water thresholds for the Responses of leaf expansion and gas exchange: A review. Field Crop Res, 47: 253 – 266

[17] Singh, B.B., O.L. Chambliss, and B. Sharma. 1997. Recent advances in cowpea breeding. Pages 30–49 in Advances in cowpea research, edited by B.B.

Farin wake	11.11(0.10)i
IT98K-497-4	8.67(0.09)i
TT 90	8.67(0.09)i
TN 5-78	8.33(0.08)i
TN 28-87	4.83(0.05)i
DNT-07	3.44(0.03)i
IT 98K 205-8	2.06(0.02)i
Dan illa	2.06(0.02)i

**

S. E. ± 19.46

Means followed by the same letters are not significantly (P<0.001) different according to Duncan's Multiple Range Test (DMRT). Means in parentheses are transformed values

TABLE 3
INCIDENCE OF SEED ROT DISEASE ON 45 COWPEA VARIETIES

Variety	Means
Dan wuri (Mai Bakin Hanci)	100(1.57)a
Jangau (Magarya)	100(1.57)a
Dan wuri (Mai Jan Hanci)	100(1.57)a
TN-121-87	95.83(1.34)ab
IT 89	86.11(1.06)bc
TN 257-87	72.22 (0.84)cd
TN-578 (Red variety)	70.83(0.79)cde
IN 92E-26	68.06(0.75)cdef
Gidimount (Jan wake)(Mirya)	51..39(0.63)cdefgh
KD-97	47.22(0.66)cdefg
IT-96D-610	47.22(0.49)defghi
Dan mora (Kwanar Dangora)	45.83(0.49)defghi
Dan misra (Rano)	45.83(0.50)defghi
Oloka (Mirya)	44.78(0.49)defghi
IT89KD-374-57	44.45(0.49)defghi
IT90K-372-1-2	41.67(0.43)defghi
KVX	41.60(0.43)defghi
TN5-78	40.28(0.42)defghi
Haladu Kadawa	35.06(0.37)defghi
IT90K-372-1-2	33.51(0.34)efghi
Dan wuri(Mai feshi)	31.94(0.34)efghi
Kyambas (Darki)	31.94(0.33)efghi
Danbathage (North)	30.56(0.31)efghi
Dan Dukku (Darki)	29.17(0.30)fg
Mai Fitila	29.17(0.30)fg
IT99k-573-1	27.78(0.28)fg
Danyagaji	24.38(0.27)fg
Oloka(Magaria NR)	23.6(0.24)gh
TN-2780	20.83(0.22)gh
TN 3-78	20.83(0.21)gh
Kanannado (Kwanar Dangora)	19.78(0.20)gh
Sama'ila	19.44(0.20)gh
Kanannado (Rano)	17.00(0.18)ghi
TN5-78	16.67(0.17)hi
KVX-100-2	16.66(0.17)hi
KVX 30-309-64	13.89(0.14)hi
TN 256-87	11.44(0.11)i

TABLE .2

LABORATORY ANALYSIS, % OF THE COLLECTED GERMLASM 2013/2014

Varieties/genotypes	Seed viability	Germination rate	Seed purity	Dead seed/infection
DAN-YAGAJI	17	42.5	100	23
TN26-6-37	7	17.5	100	33
TN5-78	11	27.5	100	29
KD-97	30	75	100	10
TN27-80	32	80	100	8
KVX	12	30	100	23
IT-89	17	42.5	100	23
IT90	33	82.5	100	7
TN-121-87	4	10	100	36
TN3-78	27	67.5	100	13
IT90K372-1-2	13	32.5	100	27
TN5-78	16	40	100	14
IT90K372-1-2	8	20	100	32
KYAMBAS	13	57.5	100	17
DAN-DUKKU	13	32.5	100	27
DANMORA	18	45	100	22
DANWURI	21	52.5	100	19
KANANNADO	29	72.5	100	11
GIDIMOUNT	10	10	100	30
IT90K-372-1-2	24	60	100	16
OLOKA	7	17.5	100	23
JANGAU	0	0	100	40
MAIFITILA	13	32.5	100	27
OLOKA	13	32.5	100	27
TN28-87	14	35	100	26
TN-378RED	5	12.5	100	35
DANMISRA	27	67.5	100	13
KANANNADO	28	70	100	12
DANWURI RED EYE	27	67.5	100	13
DANWURI BLACK EYE	17	42.5	100	23
89KD-374-57	16	40	100	24
DANILLA	12	30	100	28
KVX100-2	33	82.5	100	7
TN5-78	29	72.5	100	11
TN26-6-37	32	80	100	8
DMI-07	36	90	100	4
TN3-78	3	7.5	100	37
IN92E-26	9	22.5	100	31
KVX30-309-64	10	50	100	20
DANBATHAGE	18	45	100	22
IT99K573-1-1	10	25	100	30
IT98K097-4	28	70	100	12
IT98K205-8	35	87.5	100	5
IT96D-610	0	0	100	40

TABLE2 3
GRAIN YIELD (KG/HA)

Variety	Means
IT 98k-497-4	2668.7a
IT99k-573-1	2346.1ab
Dan dukku (Darki)	2288.5abc
TN3-78	2112.8abcd
IT96D-610	2050.4abcde
IT98K 203-8	1722.2abcdef
Dan illa	1698.4abcdef
Sama'ila	1533.7abcdefg
Mai fitila	1485.9abcdefg
IN-92E-26	1272.4abcdefg
Dan mora(Kwanar Dangora)	1270.0abcdefg

TT 90	1122.3bcdefg
IT 90K-372-1-2	1081.7bcdefg
Kyambas (Darki)	1038.3bcdefg
Haladu Kadawa	1008.5bcdefg
Dan wuri (Mai feshi)	1005.6bcdefg
KVX	990.3bcdefg
KD-97	942.0bcdefg
Oloka (Magaria NR)	922.5bcdefg
IT 89KD-374-57	820.4cdefg
DNT-07	788.6cdefg
TN5-78	745.0defg
TN28-87	742.8defg
TN256-87	694.1defg
IT 90K-372-1-2	644.3defg
TN-2780	642.8defg
TN 257-87	564.3efg
Kanannado (Rano)	543.8efg
KVX30-309-64	510.7fg
TN5-78	467.6fg
Farin wake	421.3fg
TN5-78	376.2fg
TT 89	339.5fg
Gidimount (Jan Wake)	297.3fg
TN-121-87	289.0fg
KVX-100-2	256.8fg
TN-578	208.5fg
Oloka (Mirya)	204.8fg
Danyagaji	190.3fg
Dan misra	126.3g
Danbathage (North)	109.6g
Kanannado (Kwanar Dangora)	0.0g
Dan wuri (Mai Jan Hanci)	0.0g
Jangau	0.0g
Dan wuri (Mai Bakin Hanci)	0.0g

**

S. E. \pm 753.56

Means followed by the same letters are not significantly ($P \leq 0.001$) different according to Duncan's Multiple Range Test (DMRT). Means in parentheses are transformed values