

# Interactions Study between Methyl Violet and Sodium Dodecyl Sulfate by Conductometric and Spectrophotometric Methods

Khaled Edbey, Nabil Bader, Fateh B. Eltaboni, Abdelkader Elabidi, Salah Albaba, Maher Ahmed

**Abstract**— The interactions of cationic dyes (methyl violet) with anionic surfactant sodium dodecyl sulfate (SDS) was studied in process of solubilization. The critical micelle concentration of the SDS with and without dye was determined by spectrophotometry and Specific conductometry methods. The binding constant ( $K_b$ ) and Gibbs free energy ( $\Delta G^\circ$ ) were calculated at room temperature.  $K_b$  was calculated by means of Benesi-Hildebrand Equation. Results obtained showed that the  $K_b$  and  $\Delta G^\circ$  and are found to be =  $432.77 \text{ M}^{-1}$  and  $\Delta G^\circ -15.03 \text{ KJ mol}^{-1}$  respectively. The value of  $\Delta G^\circ$  indicates that the interaction of methyl violet with micelles is spontaneous.

**Keywords**—Sodium dodecyl sulfate, CMC determination, methyl violet, dye-micelle interaction, binding constant, Gibbs free energy, conductimetry, spectrophotometry.

## I. INTRODUCTION

**S**URFACTANTS are organic compounds decrease the surface tension of water at relatively low concentrations.

Because surfactants are adsorbed mainly on the surface of the solution, creating a thin monolayer, they are called surface active agents. [1-4]

Many techniques were used for qualitative and quantitative description of dye-surfactant interactions, i.e. potentiometry [5, 6] and, conductometry [7, 8] and, or ion selective electrodes [5, 9]and. The most often used method to investigate dye-surfactant interactions is spectrophotometry. [10-14] Extensive research carried out recently has confirmed the ability of surfactants to affect the electronic absorption spectra of solutions of many dyes. the mechanism of interaction between the dye and the surfactant is still of interest.[15] In this article, a conductometric and a spectrophotometric study of the interaction of anionic surfactant (sodium dodecyl sulfate) [16-18] with cationic dye (methyl violet) [19, 20] were investigated. The aggregation behavior of SDS with dyes by spectrophotometric and conductometric methods were investigated. The critical micelle concentration of SDS with and without dye was determined. The binding constant and the Gibbs free energy were calculated.

Khaled Edbey, Nabil Bader, Fateh B. Eltaboni, Abdelkader Elabidi, Salah Albaba, Maher Ahmed, are with University of Benghazi, Libya.

## II. EXPERIMENTAL

### 2.1. Materials

Sodium dodecyl sulfate and methyl violet were obtained from Merck. A stock solution of  $2.5 \times 10^{-5} \text{ mol dm}^{-3}$  methyl violet was prepared by dissolving purified dye in distilled water. A stock solution of  $8.0 \times 10^{-2} \text{ mol dm}^{-3}$  surfactants was prepared in distilled water.



### 2.2. Procedure

The spectrophotometric measurements were recorded by a Perkin-Elmer spectrophotometer at room temperature with a matched pair of silica cuvetts having an internal thickness of 10 mm. The specific conductance measurements were carried out with Metrohm Conductometer equipped with a platinum electrode at  $25^\circ \text{C}$ . Experiments were carried out by adding different amounts of a stock surfactant solution to measuring the conductivity with and without of dye. The critical micelle concentrations were determined from the plot of specific conductivity versus concentration of the surfactant solution.

## III. RESULTS AND DISCUSSION

### 3.1 Interaction of SDS with Methyl violet by Absorption Spectroscopy:

Fig 1. Shows the structure of dye methyl violet which exists in an aqueous solution as cationic form. SDS exists in anionic form having the structure shown in (fig.2).



The effect of anionic surfactant on the absorption spectrum of methyl violet was studied at room temperature and pH 6.9. the visible spectra of aqueous methyl violet solution for several SDS concentrations ranging from  $6.4 \times 10^{-5}$  to  $1.28 \times 10^{-2}$

mol dm<sup>-3</sup> for a fixed dye concentration of 2.5×10<sup>-4</sup> mol dm<sup>-3</sup> were represents in (Fig.3). The dye exhibits a maximum absorption band at 580 nm (the band attributed to dye monomer) and a shoulder at 550 nm (the band attributed to dye dimer).

As the surfactant concentration increased gradually from 6.4×10<sup>-5</sup> to 2.56×10<sup>-4</sup> mol dm<sup>-3</sup>, below CMC, the absorbance band at 580 nm decreased (hypochromic) and the band at 550 nm diminished. The decrease in the absorbance indicates the molecular complex formation between cationic methyl violet and anionic surfactant molecules due to the electrostatic interaction. When the concentration of SDS increases (above the CMC) a significant increase in the absorbance at band 580 nm was observed and the shoulder at band 550 nm appears again. The increase in absorbance values with the increase in surfactant concentrations above CMC is regarded to be caused by the incorporation of dye molecules to micelles.

It may be thought that association of SDS anions with dye cation suppresses their mutual repulsion forces and thus favors the dye polymerization [21], but the electrostatic repulsion within the anionic moiety of SDS is reduced by the positive charge of the added dye cation. The associates in turn can further induce the formation of pre-micellar surfactant aggregate with solubilized dye content and may form other higher dye aggregate. [19, 22]




Fig.3 Visible absorption spectra of methyl violet and SDS  
3.2. Conductometric determination of the critical micelles concentration of SDS:

Another technique was employed, electrical conductivity measurements, to study the micellar aggregation behavior in aqueous solution. Figure 4 depicts the change of the specific conductance,  $\kappa$ , as a function of concentration of the surface active SDS solutions at 298 K. It can be observed that, for each surface active SDS, the results fit into two straight lines with different slopes. The break in the  $\kappa$  versus  $C$  plot originates from the micellization of amphiphilic compounds, 38 and the concentration of the break point corresponds to the cmc.

It can be seen in the figure that the slopes of the linear region above cmc are smaller than those below cmc. This is consequence of counterion binding at the surface of micellar aggregates. In other words, there is an effective loss of ionic

charges since a number of counterions are confined to the micellar surface.




Fig.4: Specific conductance of SDS plotted against the surfactant concentration.

The increasing rate of conductivity had become slower obviously after 5.9×10<sup>-3</sup>M concentration of surfactant. This is because the formation of micelle required the ionic monomers and some of the ions had been attracted towards the micelle surrounding to form the electric double layer. So the conductivity of the solution increased slower. These deviations can be explained also as a consequence of the formation of a non-conducting or a less-conducting species in solution. It is very likely that the dye cation and the surfactant anion formed a practically non-conducting, soluble ion pair. [7]

The exterior of the micelle is built up from the ionic <sup>-</sup>OSO<sub>3</sub> groups which form the Stern layer which associated by water molecules (Fig.5). The further layer that surrounding the Stern layer is composed of the positive counter ions and oriented water molecule called Gouy-Chapman layer. Both Stern layer and Gouy-Chapman layer are known as electric double layer. This double layer will maintain the stability of the colloidal system.




Fig.5: A schematic of the regions of SDS micelle and the possible site of salvation of the dye methyl violet [15].

3.2 Conductometric determination of the critical micelles concentration of SDS with dye methyl violet:

It has been observed from the experiment that the CMC of SDS surfactant start to form at low concentration (Fig.6). The

low value of the CMC results as the electrostatic repulsion within the anionic moiety of SDS is reduced by positive charge of added dye anion. [16].




Fig.6: Specific conductance of SDS with dye plotted against the surfactant concentration.

### 3.3 Spectrophotometric determination of the critical micelles concentration of SDS with dye methyl violet:

CMC was determined experimentally for SDS in range from  $6.4 \times 10^{-5}$  M to  $1.6 \times 10^{-2}$  M. (Fig.7) shows determination of the CMC of SDS using spectrophotometric method. The CMC value was found to be 0.0048 M.





Fig.7: Determination of CMC of SDS by spectrophotometer

### 3.3 Determination of binding constant of SDS ( $K_b$ ):

The interaction between the dye and micelles can be described as:



Where D is the dye, M the micelle, DM the dye-micelle complexes and  $K_b$  is the binding constant. Benesi -hildebrand equation gives more precise parameters as binding constant  $K_b$ .

[19, 23]

$$\frac{D_T}{\Delta A} = \frac{1}{(\epsilon_m - \epsilon_0)} + \frac{1}{kb(\epsilon_m - \epsilon_0)C_m}$$

Where  $D_T$  is the concentration of dye,  $\Delta A = A - A_0$  is the difference between the absorbance of dye in the presence and absence of surfactant,  $\epsilon_m$  is the molar extinction coefficient of dye fully bound to micelles,  $\epsilon_0$  is the molar extinction coefficient of the methyl violet,  $K_b$  is the binding constant,  $C_m$  is the Concentration of the micellized surfactant.

$$C_m = C_s - CMC$$




Fig. 8. The plot of  $DT / \Delta A$  against  $1/C_m$  for methyl violet in SDS

Results obtained from the spectral measurements (fig. 8) showed that the binding constant  $K_b$  is found to be  $= 432.77 \text{ M}^{-1}$ .

### 3.4 Determination of standard free energy change:

The thermodynamic parameter  $\Delta G^\circ$  which is an indicator of the tendency of binding of methyl violet to micelles was calculated using the following equation:

$$\Delta G^\circ = -RT \ln K_b$$

Where  $\Delta G^\circ$  is the standard free energy change, R is the universal gas constant and T is the room temperature.

The value of  $\Delta G^\circ$  is found to be  $= -15.03 \text{ KJ mol}^{-1}$  which indicates that the interaction of methyl violet with micelle is spontaneous.

## IV. CONCLUSION

Based on the study of the interaction of methyl violet with SDS, the following conclusions can be drawn:

The effect of the dye on the micellization of SDS in aqueous media was successfully analyzed using electrical conductivity and UV-vis measurements. Critical micelles concentration of SDS with and without dye was determined by conductimetric and spectrophotometric methods. The binding constant ( $K_b$ ), Gibbs free energy parameter ( $\Delta G^\circ$ ) were calculated. The aggregation behavior of SDS with dye was studied by absorption spectroscopy).

REFERENCES

- [1] B. Cai, X. Li, Y. Yang, et al., Surface properties of Gemini surfactants with pyrrolidinium head groups. *J Colloid Interface Sci*, Vol. 370(1), pp. 111-6.  
<http://dx.doi.org/10.1016/j.jcis.2011.12.025>
- [2] H. Chen, J. Zhao, J. Wu, et al., Isotherm, thermodynamic, kinetics and adsorption mechanism studies of methyl orange by surfactant modified silkworm exuviae. *J Hazard Mater*, Vol. 192(1), pp. 246-54.  
<http://dx.doi.org/10.1016/j.jhazmat.2011.05.014>
- [3] S. Ghosh and J. Dey, Interaction of sodium N-lauroylsarcosinate with N-alkylpyridinium chloride surfactants: spontaneous formation of pH-responsive, stable vesicles in aqueous mixtures. *J Colloid Interface Sci*, Vol. 358(1), pp. 208-16.  
<http://dx.doi.org/10.1016/j.jcis.2011.02.054>
- [4] M. J. Rosen and J. T. Kunjappu, *Micelle Formation by Surfactants, in Surfactants and Interfacial Phenomena*. 2012, John Wiley & Sons, Inc. p. 123-201.  
<http://dx.doi.org/10.1002/9781118228920.ch3>
- [5] M. Kert and B. Simončič, The influence of nonionic surfactant structure on the thermodynamics of anionic dye-cationic surfactant interactions in ternary mixtures. *Dyes and Pigments*, Vol. 79(1), pp. 59-68  
<http://dx.doi.org/10.1016/j.dyepig.2008.01.005>
- [6] B. M. Razavizadeh, M. Mousavi-Khoshdel, H. Gharibi, et al., Thermodynamic studies of mixed ionic/nonionic surfactant systems. *J Colloid Interface Sci*, Vol. 276(1), pp. 197-207.  
<http://dx.doi.org/10.1016/j.jcis.2004.03.036>
- [7] S. Bračko and J. Špan, Conductometric investigation of dye-surfactant ion pair formation in aqueous solution. *Dyes and Pigments*, Vol. 45(2), pp. 97-102  
[http://dx.doi.org/10.1016/S0143-7208\(00\)00016-4](http://dx.doi.org/10.1016/S0143-7208(00)00016-4)
- [8] T. Joshi, J. Mata, and T. Patel, Conductometry and Thermodynamics Study of Metal Dodecyl Sulfates in Aqueous Solution. *Journal of Dispersion Science and Technology*, Vol. 28(8), pp. 1158-1163  
<http://dx.doi.org/10.1080/01932690701525510>
- [9] J. L. Lopez-Fontan, A. Gonzalez-Perez, J. Costa, et al., The critical micelle concentration of tetraethylammonium perfluorooctylsulfonate in water. *J Colloid Interface Sci*, Vol. 294(2), pp. 458-65.  
<http://dx.doi.org/10.1016/j.jcis.2005.07.029>
- [10] H. Akbas and C. Kartal, C.I Reactive Orange 16-dodecylpyridinium chloride interactions in electrolytic solutions. *Spectrochim Acta A Mol Biomol Spectrosc*, Vol. 65(1), pp. 95-9.  
<http://dx.doi.org/10.1016/j.saa.2005.09.033>
- [11] B. Arıkan and M. Tunçay, The effect of SDS micelles on reduction of toluidine blue by ascorbic acid in acid medium. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, Vol. 273(1-3), pp. 202-207.  
<http://dx.doi.org/10.1016/j.colsurfa.2005.08.025>
- [12] B. S. Berlett, R. L. Levine, and E. R. Stadtman, Use of isosbestic point wavelength shifts to estimate the fraction of a precursor that is converted to a given product. *Anal Biochem*, Vol. 287(2), pp. 329-33.  
<http://dx.doi.org/10.1006/abio.2000.4876>
- [13] R. Hosseinzadeh, R. Maleki, A. A. Matin, et al., Spectrophotometric study of anionic azo-dye light yellow (X6G) interaction with surfactants and its micellar solubilization in cationic surfactant micelles. *Spectrochim Acta A Mol Biomol Spectrosc*, Vol. 69(4), pp. 1183-7.  
<http://dx.doi.org/10.1016/j.saa.2007.06.022>
- [14] B. Simončič and F. Kovac, A study of dye-surfactant interactions. Part 2. The effect of purity of a commercial cationic azo dye on dye-surfactant complex formation. *Dyes and Pigments*, Vol. 40(1), pp. 1-9.  
[http://dx.doi.org/10.1016/S0143-7208\(98\)00028-X](http://dx.doi.org/10.1016/S0143-7208(98)00028-X)
- [15] M. E. D. Garcia and A. Sanz-Medel, Dye-surfactant interactions: a review. *Talanta*, Vol. 33(3), pp. 255-264  
[http://dx.doi.org/10.1016/0039-9140\(86\)80060-1](http://dx.doi.org/10.1016/0039-9140(86)80060-1)
- [16] Benito, M. A. García, C. Monge, et al., Spectrophotometric and conductimetric determination of the critical micellar concentration of sodium dodecyl sulfate and cetyltrimethylammonium bromide micellar systems modified by alcohols and salts. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, Vol. 125(2-3), pp. 221-224.  
[http://dx.doi.org/10.1016/S0927-7757\(97\)00014-9](http://dx.doi.org/10.1016/S0927-7757(97)00014-9)
- [17] D. Das and K. Ismail, Aggregation and adsorption properties of sodium dodecyl sulfate in water-acetamide mixtures. *J Colloid Interface Sci*, Vol. 327(1), pp. 198-203.  
<http://dx.doi.org/10.1016/j.jcis.2008.07.045>
- [18] R. K. Dutta, R. Chowdhury, and S. N. Bhat, Effect of association of sulfonephthalein dyes with sodium dodecyl sulfate micelles on their acid-base equilibria. *Journal of the Chemical Society, Faraday Transactions*, Vol. 91(4), pp. 681-686.  
<http://dx.doi.org/10.1039/ft9959100681>
- [19] M. Sarkar and S. Poddar, Spectral studies of methyl violet in aqueous solutions of different surfactants in supermicellar concentration region. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, Vol. 55(9), pp. 1737-1742.  
[http://dx.doi.org/10.1016/S1386-1425\(98\)00344-8](http://dx.doi.org/10.1016/S1386-1425(98)00344-8)
- [20] G. Vinitha and A. Ramalingam, Spectral characteristics and nonlinear studies of methyl violet 2B dye in liquid and solid media. *Laser Physics*, Vol. 18(1), pp. 37-42.  
<http://dx.doi.org/10.1134/S1054660X08010076>
- [21] S. P. Moulik, S. Ghosh, and A. R. Das, Interaction of acridine orange monohydrochloride dye with sodiumdodecylsulfate, (SDS) cetyltrimethylammoniumbromide (CTAB) and p-tert-octylphenoxypolyoxy ethanol (Triton X 100) surfactants. *Colloid and Polymer Science*, Vol. 257(6), pp. 645-655.  
<http://dx.doi.org/10.1007/BF01548834>
- [22] M. Sarkar and S. Poddar, Studies on the Interaction of Surfactants with Cationic Dye by Absorption Spectroscopy. *J Colloid Interface Sci*, Vol. 221(2), pp. 181-185.  
<http://dx.doi.org/10.1006/jcis.1999.6573>
- [23] S. Göktürk and M. Tunçay, Spectral studies of safranin-O in different surfactant solutions. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, Vol. 59(8), pp. 1857-1866.  
[http://dx.doi.org/10.1016/S1386-1425\(02\)00418-3](http://dx.doi.org/10.1016/S1386-1425(02)00418-3)